



SWEET PEPPERS FOR THE EAST COAST

- SE Southeast
- MW Midwest
- NE Northeast
- HT High Tunnel

VARIETY	TYPE	REGION/ GROWING METHOD	RELATIVE MATURITY*	FRUIT SIZE/SHAPE	FRUIT COLOR	PLANT HABIT	DISEASE RESISTANCE	USES & REMARKS
Big Stack S10®	Bell	● MW ● NE	Early	XL, Jumbo/Blocky		Moderate	HR: Tm: 0, IR: Pc / X spp 0-10	Exceptional level of Pc tolerance and very early maturing
NEW Cashout S10®	Bell	● SE ● MW ● NE	Early	Large, XL/Blocky		Moderate, good cover	HR: PVY: 0, 1 / Tm: 0, IR: Pc / TSWV / X spp: 0-10	Robust disease package including very high Pc tolerance. Early set with the flexibility of extended harvest
High Roller S10®	Bell	● SE	Mid-early	Large, XL/Blocky		Vigorous	HR: Tm: 0, IR: TSWV / X spp: 0-10	Continuous setting under cool or hot conditions
Mercer	Bell	● SE ● MW ● NE ● HT	Mid-early	XL/Blocky		Moderate, good cover	HR: Tm: 0 / X spp: 0-3, 7, 8, IR: Pc	Well adapted for the Northeast where Phytophthora capsici can be a problem. Classy fruit
Nitro S10®	Bell	● SE ● MW	Mid	Large, XL/Blocky		Moderate, excellent cover	HR: Tm: 0, IR: Pc / TSWV: 0 / X spp: 0-10	Heat tolerant with a broad disease package and place-pack quality
NEW Payline S10®	Bell	● SE	Mid	XL/Blocky		Vigorous, good cover	HR: Tm: 0 // IR: TSWV / X spp: 0-10	Extended harvest of high quality fruit. Focused for Winter and spring production slots
Takara	Shishito	● SE ● MW ● NE ● HT	Very early	Small/Elongated		Compact, well branched		Takara is a classic Japanese Shishito type pepper that have small thin walled fruit, typically harvested at mature green
Turbo S10®	Bell	● SE	Early	Large, XL/Blocky		Compact, excellent cover	HR: Tm: 0, IR: TSWV / X spp: 0-10	Heat tolerant. Very early with dark green fruit
Lemon Delite	Snacking	● SE ● MW ● NE ● HT	Very early	Medium, large/Flat, conical		Compact, good cover, medium, tall, vigorous, short internodes	HR: Tm: 0 / X spp: 0-3, 7, 8	Lemon Delite has a mid-strong vigorous plant with good cover & continuous setting ability; fresh market
NEW Golden Delite S10®	Snacking	● SE ● MW ● NE ● HT	Early	Large, XL/Elongated		Moderate plants with good cover, extended harvest types	HR: Tm: 0 IR: TSWV: 0 / X spp: 0-10	Golden Delite S10® is widely adapted, with a strong disease package for crop security & ability to set in many climates; fresh market
Orange Delite	Snacking	● SE ● MW ● NE ● HT	Very early	Medium, large/Flat, conical		Compact, good cover, medium, tall, vigorous, short internodes	HR: Tm: 0 / X spp: 0-3, 7, 8	Orange Delite has a mid-strong vigorous plant with good cover & continuous setting ability; fresh market
NEW Amber Delite S10®	Snacking	● SE ● MW ● NE ● HT	Early	Large, XL/Elongated		Moderate plants with good cover, extended harvest types	HR: Tm: 0 IR: TSWV: 0 / X spp: 0-10	Amber Delite S10® is widely adapted, with a strong disease package for crop security & ability to set in many climates; fresh market
Ruby Delite	Snacking	● SE ● MW ● NE ● HT	Very early	Medium, large/Flat, conical		Compact, good cover, medium, tall, vigorous, short internodes	HR: Tm: 0 / X spp: 0-3, 7, 8	Ruby Delite has a mid-strong vigorous plant with good cover & continuous setting ability; fresh market
NEW Crimson Delite S10®	Snacking	● SE ● MW ● NE ● HT	Early	Large, XL/Elongated		Moderate plants with good cover, extended harvest types	HR: Tm: 0 IR: TSWV: 0 / X spp: 0-10	Crimson Delite S10® is widely adapted, with a strong disease package for crop security & ability to set in many climates; fresh market

Disease Terminology: HR = High Resistance; IR = Intermediate Resistance **Disease Abbreviation Code:** Lt - Powdery mildew | Mi - Root-knot | Pc - Phytophthora crown and root rot | PVY - Potato virus Y | Tm - Tobamovirus | TSWV - Tomato spotted wilt | X spp - Bacterial spot
 * Relative Maturity are an approximation and may fluctuate due to varying planting times, location and condition

S10® - Signifies intermediate resistance to all 10 races of bacterial leaf spot in pepper. Bacterial spot of tomato and pepper is caused by four bacterial species (Xanthomonas euvesicatoria, X. vesicatoria, X. perforans, and X. gardneri). Of the four species, X. euvesicatoria appears to be the main causal agent for bacterial spot of pepper. At this time, 10 races of the pathogen have been identified on pepper. Because of the difficulty in differentiating the bacterial species with non-DNA based methods, the seed industry currently recognizes the causal agent by its former scientific name of Xanthomonas capestris pv. vesicatoria prior to its reclassification.

DISCLAIMER: Any representations and other disclosed information are based on our observations and/or information from other sources. Crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, the environment, including management, and other uncontrollable factors that may alter expected performance. Triploid hybrid watermelon varieties will on occasion produce white and hard, dark vestigial seed coats and, therefore, are not warranted to be completely "seedless". Statements on the reaction of varieties to a specific pathogen, pest or stress are based on evaluation under defined conditions. These reactions can be affected by changes in environmental and biological factors, especially new pathogen races, pest biotypes or vectors of disease agents. Therefore, Sakata gives no warranty, express or implied, for crop performance relative to the information given; nor does Sakata accept any liability for any loss, direct, indirect, or consequential, that may arise from any cause. Please read all seed package labeling carefully as it contains terms and conditions of sale.